


Solubility Rules for some ionic compounds in water**Soluble Ionic Compounds**

1. All sodium (Na^+), potassium (K^+), and ammonium (NH_4^+) salts are SOLUBLE.
2. All nitrate (NO_3^-), acetate (CH_3CO_2^-), chlorate (ClO_3^-), and perchlorate (ClO_4^-) salts are SOLUBLE.
3. All chloride (Cl^-), bromide (Br^-), and iodide (I^-) salts are SOLUBLE -- EXCEPT those also containing: lead, silver, or mercury (I) (Pb^{2+} , Ag^+ , Hg_2^{2+}) which are NOT soluble.
4. All sulfate (SO_4^{2-}) salts are SOLUBLE -- EXCEPT those also containing: calcium, silver, mercury (I), strontium, barium, or lead (Ca^{2+} , Ag^+ , Hg_2^{2+} , Sr^{2+} , Ba^{2+} , Pb^{2+}) which are NOT soluble.



Not Soluble Ionic Compounds

5. Hydroxide (OH^-) and oxide (O^{2-}) compounds are NOT SOLUBLE -- EXCEPT those also containing: sodium, potassium, or barium (Na^+ , K^+ , Ba^{2+}) which are soluble.
6. Sulfide (S^{2-}) salts are NOT SOLUBLE -- EXCEPT those also containing: sodium, potassium, ammonium, or barium (Na^+ , K^+ , NH_4^+ , Ba^{2+}) which are soluble.
7. Carbonate (CO_3^{2-}) and phosphate (PO_4^{3-}) salts are NOT SOLUBLE -- EXCEPT those also containing: sodium, potassium, or ammonium (Na^+ , K^+ , NH_4^+), which are soluble.


Some common ions:**Bond Dissociation Energies (kJ mol^{-1}) (gas phase)**

Bond	D	Bond	D	Bond	D
H-H	436	C-C	346	N-N	163
C-H	413	C=C	610	N=N	418
$\text{C}\equiv\text{O}$	1046	$\text{C}\equiv\text{N}$	887	$\text{N}\equiv\text{N}$	945
N-H	391	O-O	146	C-O	358
O-H	463	O=O	498	C=O	745
C-F	485	F-F	155	N-F	283
C-Cl	339	Cl-Cl	242	N-Cl	192
C-I	213	I-I	151	N-I	169

5 Given the following information:



what is the standard enthalpy change for the reaction:



- 1) 155 kJ mol^{-1} 2) -146 kJ mol^{-1} 3) 9.2 kJ mol^{-1}
4) 146 kJ mol^{-1} 5) not enough information to determine




6 The root mean square speed of molecules in a sample of F_2 gas is 890 m/s. What is the temperature of the gas?

- 1) 513 K 2) 890 K 3) 127 K 4) 1208 K 5) 233 K

7 A 2.38 mol sample of He gas is confined in a 62.5 liter container at 62.5 °C. If 1.28 mol of F_2 gas is added while maintaining constant temperature, the average kinetic energy per molecule will:

- 1) decrease 2) remain the same 3) increase
4) not enough information 5) I don't have a clue

8. Which listing below correctly orders the molecules by increasing root mean square molecular speed (slowest \rightarrow fastest)?



9. A sample of Cl_2 gas is confined in a 2.0 liter container at 50°C . Then 2.5 mol of He is added, holding both the volume and temperature constant. The pressure will increase because:

1) With more molecules per unit volume, there are more molecules hitting the walls of the container.

2) As the number of molecule-wall collisions increases, the force per collision increases.

3) With more molecules in the container, the molecules have higher average speeds.

4) With higher average speeds, on average the molecules hit the walls of the container with more force.

5) None of the Above

10. What is the average kinetic energy of an N_2 molecule confined in 3.1 L at 1.0 atm and 25°C ?

1) $5.71 \times 10^3 \text{ J}$


2) $9.48 \times 10^3 \text{ J}$

3) $6.17 \times 10^{-21} \text{ J}$

4) $5.71 \times 10^{-21} \text{ J}$

5) $3.21 \times 10^{-21} \text{ J}$

Consider the molecular orbital energy diagram shown at right.




- 11 The energy level denoted “b” refers to:
- 1) a bonding molecular orbital
 - 2) an antibonding molecular orbital
 - 3) a nonbonding molecular orbital
 - 4) an atomic orbital
- 12 The electrons in the orbital represented by energy level “b”:
- 1) are distributed more toward X
 - 2) are distributed more toward Y
 - 3) are equally distributed between X and Y
- 13 The molecule XY is the diatomic $(\text{He-H})^+$. What is its bond order?
- 1) 0.0
 - 2) 0.5
 - 3) 1.0
 - 4) 1.5
 - 5) 2.0
14. What is the energy of ultraviolet light with frequency 1.36×10^{15} Hz?
- 1) 126 kJ mol^{-1}
 - 2) 196 kJ mol^{-1}
 - 3) 427 kJ mol^{-1}
 - 4) 544 kJ mol^{-1}
 - 5) 832 kJ mol^{-1}
15. Consider two cases for emission from the hydrogen atom:
- | | |
|---|---|
| Case 1:
Electron goes from $n=4$ to $n=3$ | Case 2:
Electron goes from $n=6$ to $n=2$ |
|---|---|
- Compare the energies of the photons emitted:
- 1) $E_{\text{case 1}} > E_{\text{case 2}}$
 - 2) $E_{\text{case 1}} < E_{\text{case 2}}$
 - 3) $E_{\text{case 1}} = E_{\text{case 2}}$

16. Consider the energy vs temperature diagram at right, describing the transitions of water from ice to steam:

The segment labeled (d) is described best with which parameter below:

- 1) $\Delta H^\circ_{\text{fus}}$ 2) $\Delta H^\circ_{\text{vap}}$ 3) C_{ice}
 4) C_{liquid} 5) C_{steam}



17. The following information is given for bismuth, Bi, at 1atm:

$$\text{boiling pt} = 1627^\circ\text{C} \quad H_{\text{vap}}^{1627^\circ\text{C}, 1\text{atm}} = 172 \text{ kJ mol}^{-1} \quad C_{\text{liquid Bi}} = 0.151 \text{ J g}^{-1} \text{ K}^{-1}$$

$$\text{melting pt} = 271^\circ\text{C} \quad H_{\text{fus}}^{271^\circ\text{C}, 1\text{atm}} = 11.0 \text{ kJ mol}^{-1} \quad C_{\text{solid Bi}} = 0.126 \text{ J g}^{-1} \text{ K}^{-1}$$

At a pressure of 1 atm, what amount of heat is needed to melt a 35.2 g sample of solid bismuth at its normal melting point of 271 °C?

- 1) 4.21 kJ 2) 13.8 kJ 3) 0.561 kJ 4) 9.67 kJ 5) 1.85 kJ
18. At a pressure of 1 atm, what amount of heat is needed to take a 35.2 g sample of bismuth from 200°C to 400°C?
- 1) 2.85 kJ 2) 15.4 kJ 3) 32.6 kJ 4) 9.67 kJ 5) 14.3 kJ

19. Which ion has the smallest radius?

- 1) K^+ 2) Ca^{2+} 3) P^{3-} 4) S^{2-} 5) all the same

20. Consider the following samples:

- a) 0.531 moles of CH_4 in a 6.18 L container at a temperature of 308K
b) 0.281 moles of CH_4 in a 2.77 L container at a temperature of 388K
c) 0.569 moles of CH_4 in a 1.42 L container at a temperature of 453K
d) 0.212 moles of CH_4 in a 5.95 L container at a temperature of 298K

Which has the lowest average molecular speed?

- 1) a 2) b 3) c 4) d 5) all the same

21. HNO_3 is (data at the front of the exam provide a clue):

- 1) a strong base 2) a weak acid 3) a weak base
4) a strong acid 5) none of the above

22. Reactions in water that produce gases tend to be:


- 1) favorable 2) ugly 3) unfavorable
4) exothermic 5) endothermic

23. Which reaction below is a redox reaction?

- 1) $NaOH(aq) + HNO_3(aq) \rightarrow NaNO_3(aq) + H_2O(l)$
2) $Na_2CO_3(aq) + 2HClO_4(aq) \rightarrow CO_2(g) + H_2O(l) + 2NaClO_4$
3) $CdCl_2(aq) + Na_2S(aq) \rightarrow CdS(s) + 2NaCl(aq)$
4) $Zn(OH)_2(s) + H_2SO_4(aq) \rightarrow ZnSO_4(aq) + 2H_2O(l)$
5) None of the above

24. The net ionic equation for the reaction of zinc sulfate and sodium hydroxide is:

- 1) $Zn^{2+}(aq) + 2OH^-(aq) \rightarrow Zn(OH)_2(s) + Na_2SO_4(aq)$
2) $ZnSO_4(aq) + 2NaOH(aq) \rightarrow Zn(OH)_2(aq) + Na_2SO_4(aq)$
3) $Zn^{2+}(aq) + 2OH^-(aq) \rightarrow Zn(OH)_2(s)$
4) $Zn^{2+}(aq) + 2OH^-(aq) \rightarrow Zn(OH)_2(aq)$
5) No *net* reaction occurs

25. Which element has the lowest ionization energy?
1) In 2) Ga 3) Tl 4) B 5) all the same
26. Draw the Lewis structure for CO^{2-} . What is the hybridization on oxygen?
1) sp^3d 2) sp^4 3) sp^3 4) sp^2 5) sp
27. Draw the Lewis structure for XeOF_4 (Xe is the central atom). What is the hybridization on **Xe**?
1) sp^3d^2 2) sp^3d^3 3) sp^3d 4) sp^3 5) sp^2
28. The molecule XeOF_4 is:
1) polar 2) nonpolar 3) can't tell
29. The correct molecular formula for the molecule at right is:
1) $\text{C}_2\text{O}_2\text{H}_4$ 2) CO_2H_4 3) C_2OH_4 4) $\text{C}_2\text{O}_2\text{H}_3$
- 
30. A specific isotope of an ion from a given element has 8 protons, 7 neutrons, and 10 electrons. The ion is:
1) O^{2-} 2) Ne^{3-} 3) P^{3-} 4) N^{3-} 5) Mn^{3+}
31. What is the formula of the ionic compound formed in the reaction of elemental **Mg** and **O₂**?
1) MgO 2) Mg_2O 3) Mg_2O_3 4) Mg_3O_2 5) MgO_2
32. What is the (mass) percent composition of **C** in **C₄H₈**?
1) 85.6% 2) 14.4% 3) 50.0% 4) 88.3% 5) 11.7%

33. What is the wavelength of ultraviolet light with frequency 1.18×10^{15} Hz?
1) 209 nm 2) 254 nm 3) 280 nm 4) 190 nm 5) 350 nm
34. What is the maximum number of orbitals that can be identified by the set of quantum numbers $n=+5$ $l=+3$?
1) 2 2) 3 3) 5 4) 6 5) 7
35. Consider the molecule ClF_4^- How many lone **pairs** are on the central atom?
1) 1 2) 2 3) 3 4) 6 5) 0
36. Light is given off by a sodium or mercury containing street light when the atoms are excited. The light you see arises for which of the following reasons?
1) Electrons are moving from a given energy level to one of lower n
2) Electrons are moving from a given energy level to one of higher n
3) Electrons are being removed from the atom, thereby creating a metal cation
37. Consider the molecule ClF_4^- What is the electron pair geometry?
1) Trigonal bipyramidal 2) Octahedral 3) linear
4) Trigonal planer 5) Tetrahedral

38. Which of the following has the highest affinity for electrons?
1) B 2) N 3) As 4) P 5) Ge
39. In ionizing elemental lithium to Li^+ , from which orbital is an electron removed?
1) 1s 2) 2s 3) 3s 4) 2p 5) 3p
40. In the molecule **formaldehyde** CH_2O , what is the approximate HCO bond angle?
1) 180° 2) 90° 3) 109° 4) 120° 5) 60°
41. If you completely react 0.678 g of iodine (I_2), what mass of NI_3 can be produced?
1) 0.276 g 2) 0.678 g 3) 0.226 g 4) 0.876 g 5) 0.351 g

- 42 Nitrogen triiodide (NI_3) is unstable, reacting to form N_2 (g) and I_2 (g), and evolving heat.



Spontaneous decomposition of 0.5 g of NI_3 (s) produces what volume of gas at 200°C and 1 atm pressure?

- 1) 28.7 L 2) 0.197 L 3) 0.098 L 4) 14.4 L 5) 0.731 L

- 43 Using the Table of Bond Dissociation Energies at the front of the exam, predict ΔH° for the spontaneous decomposition of nitrogen triiodide above.

- 1) -256 kJ mol^{-1} 2) -927 kJ mol^{-1} 3) -384 kJ mol^{-1}
4) -35 kJ mol^{-1} 5) $+927 \text{ kJ mol}^{-1}$

- 44 What is the molecular geometry of nitrogen triiodide?

- 1) tetrahedral 2) square planar 3) trigonal planar
4) octahedral 5) trigonal pyramidal


- 45 What is the hybridization on N in nitrogen triiodide?

- 1) sp 2) sp^2 3) sp^3 4) sp^4 5) $\text{sp}^3 \text{d}$

46 Which do you expect to have the longest bond length?

- 1) NI_3 2) NBr_3 3) NCl_3 4) NF_3 5) can't tell

47 In class, we saw the following reaction (unbalanced).



In the correctly balanced reaction, what is the stoichiometry coefficient preceding Br_2 (all coefficients should be integral)?

- 1) 1 2) 2 3) 3 4) 4 5) 6

48 In the reaction above of aluminum and bromine, which is the oxidizing agent?

- 1) Al (s) 2) $\text{Br}_2 \text{ (l)}$

49 What is the electron pair geometry in AlBr_3 ?

- 1) tetrahedral 2) square planar 3) trigonal planar
4) octahedral 5) trigonal pyramidal

50 What is the catalog number for this class?

- 1) 123 2) 111 3) 86 4) 345 5) 68.6 g