


Solubility Rules for some ionic compounds in water**Soluble Ionic Compounds**

1. All sodium (Na^+), potassium (K^+), and ammonium (NH_4^+) salts are SOLUBLE.
2. All nitrate (NO_3^-), acetate (CH_3CO_2^-), chlorate (ClO_3^-), and perchlorate (ClO_4^-) salts are SOLUBLE.
3. All chloride (Cl^-), bromide (Br^-), and iodide (I^-) salts are SOLUBLE -- EXCEPT those also containing: lead, silver, or mercury (I) (Pb^{2+} , Ag^+ , Hg_2^{2+}) which are NOT soluble.
4. All sulfate (SO_4^{2-}) salts are SOLUBLE -- EXCEPT those also containing: calcium, silver, mercury (I), strontium, barium, or lead (Ca^{2+} , Ag^+ , Hg_2^{2+} , Sr^{2+} , Ba^{2+} , Pb^{2+}) which are NOT soluble.

Not Soluble Ionic Compounds

5. Hydroxide (OH^-) and oxide (O^{2-}) compounds are NOT SOLUBLE -- EXCEPT those also containing: sodium, potassium, or barium (Na^+ , K^+ , Ba^{2+}) which are soluble.
6. Sulfide (S^{2-}) salts are NOT SOLUBLE -- EXCEPT those also containing: sodium, potassium, ammonium, or barium (Na^+ , K^+ , NH_4^+ , Ba^{2+}) which are soluble.
7. Carbonate (CO_3^{2-}) and phosphate (PO_4^{3-}) salts are NOT SOLUBLE -- EXCEPT those also containing: sodium, potassium, or ammonium (Na^+ , K^+ , NH_4^+), which are soluble.

Some common ions:**Bond Dissociation Energies (kJ mol^{-1}) (gas phase)**

| Bond | D | Bond | D | Bond | D |
|--------------------------|------|--------------------------|-----|--------------------------|-----|
| H-H | 436 | C-C | 346 | N-N | 163 |
| C-H | 413 | C=C | 610 | N=N | 418 |
| $\text{C}\equiv\text{O}$ | 1046 | $\text{C}\equiv\text{N}$ | 887 | $\text{N}\equiv\text{N}$ | 945 |
| N-H | 391 | O-O | 146 | C-O | 358 |
| O-H | 463 | O=O | 498 | C=O | 745 |
| C-F | 485 | F-F | 155 | N-F | 283 |
| C-Cl | 339 | Cl-Cl | 242 | N-Cl | 192 |
| C-I | 213 | I-I | 151 | N-I | 169 |

1. In an exothermic process:

- 1) work is performed on the surroundings
- 2) heat is transferred to the surroundings
- 3) work is performed on the system
- 4) heat is transferred to the system

(2)**Ch 6.1 – Energy: basic principles**2. A positive value of ΔE means that:

- 1) heat is transferred to the surroundings
- 2) heat is transferred to the system
- 3) energy in the form of heat and/or work is transferred to the surroundings
- 4) energy in the form of heat and/or work is transferred to the system

(4)**Ch 6.4 – Energy: 1st Law of Thermo**3. An automobile engine generates **2757** Joules of heat that must be carried away by the cooling system. The internal energy changes by **-3852** Joules in this process.

How much work to push the pistons is available in this process?

- 1) 4918 J
- 2) 1095 J
- 3) 683 J
- 4) 6283 J
- 5) 1277 J

$$\Delta E = q + w \quad w = \Delta E - q = (-3852 \text{ J}) - (-2757 \text{ J}) = -1095 \text{ J}$$

(2) w is negative. The system does work on the surroundings.**Ch 6.4 – Energy: 1st Law of Thermo**4. What is the minimum amount of ice at 0°C that must be added to the contents of a can of diet cola (340 mL) to cool it from 20°C to 5°C ? Assume that diet cola has the properties of pure water ($\Delta H_{\text{fusion}}^{\text{water}} = 333 \text{ J g}^{-1}$ $d_{\text{water}} = 1.0 \text{ g mL}^{-1}$)

- 1) 34.2 g
- 2) 64.1 g
- 3) 87.6 g
- 4) 10.2 g
- 5) 125 g

$$m_{\text{cola}} = (340 \text{ mL})(1.00 \text{ g mL}^{-1}) = 340 \text{ g} \quad m_{\text{ice}} = ?$$

$$\Delta T = [(20 + 273) - (5 + 273)] \text{ K} = -15 \text{ K}$$

(2) $\Delta H_{\text{fusion}}^{\text{water}} = 333 \text{ J g}^{-1}$

$$q_{\text{cola}} + q_{\text{ice}} = 0 = [C_{\text{cola}} m_{\text{cola}} \Delta T] + [m_{\text{ice}} \Delta H_{\text{fusion}}^{\text{water}}]$$



$$m_{\text{ice}} \Delta H_{\text{fusion}}^{\text{water}} = -C_{\text{cola}} m_{\text{cola}} \Delta T$$

$$m_{\text{ice}} = \frac{-C_{\text{cola}} m_{\text{cola}} \Delta T}{\Delta H_{\text{fusion}}^{\text{water}}} = \frac{-(4.184 \text{ J g}^{-1})(340 \text{ g})(-15 \text{ K})}{333 \text{ J g}^{-1}} = 64.1 \text{ g}$$


(5)

Ch 6.3 – Specific heat capacity and phase change, see Example 6.4 on page 249


5. Given the following information:



what is the standard enthalpy change for the reaction:



- 1) 155 kJ mol^{-1} 2) -146 kJ mol^{-1} 3) 9.2 kJ mol^{-1}
 4) 146 kJ mol^{-1} 5) not enough information to determine

**Ch 6.7 – Hess's Law (Owl, Unit 6-6c)**

6. The root mean square speed of molecules in a sample of F_2 gas is 890 m/s. What is the temperature of the gas?

- 1) 513 K 2) 890 K 3) 127 K 4) 1208 K 5) 233 K

$$\overline{u^2} = \frac{3RT}{M}$$

$$(4) \quad T = \frac{M\sqrt{\overline{u^2}}^2}{3R} = \frac{38.0 \text{ g mol}^{-1} (890 \text{ m s}^{-1})^2}{3(8.314 \text{ J K}^{-1} \text{ mol}^{-1})} \frac{\text{J}}{\text{kg m}^2 \text{ s}^{-2}} \frac{\text{kg}}{10^3 \text{ g}} = 1208 \text{ K}$$

Ch 12.6 – Kinetic theory of gases

7. A 2.38 mol sample of He gas is confined in a 62.5 liter container at 62.5 °C. If 1.28 mol of F_2 gas is added while maintaining constant temperature, the average kinetic energy per molecule will:

- 1) decrease 2) remain the same 3) increase
 4) not enough information 5) I don't have a clue



Ch 12.6 – Kinetic theory of gases

8b. Which listing below correctly orders the molecules by increasing root mean square molecular speed (slowest → fastest)?

1) $\text{CO}_2 < \text{Ar} < \text{N}_2 < \text{H}_2$

2) $\text{Ar} < \text{CO}_2 < \text{N}_2 < \text{H}_2$

3) $\text{H}_2 < \text{N}_2 < \text{CO}_2 < \text{Ar}$

4) $\text{H}_2 < \text{N}_2 < \text{Ar} < \text{CO}_2$

(1) $\sqrt{u^2} = \sqrt{\frac{3RT}{M}}$ **Molar masses: 48 > 40 > 28 > 2 (OWL 12-x)**

Ch 12.6 – kinetic theory, rms speed and molar mass.

9. A sample of Cl_2 gas is confined in a 2.0 liter container at 50 °C. Then 2.5 mol of He is added, holding both the volume and temperature constant. The pressure will increase because:

1) With more molecules per unit volume, there are more molecules hitting the walls of the container.

2) As the number of molecule-wall collisions increases, the force per collision increases.

3) With more molecules in the container, the molecules have higher average speeds.

4) With higher average speeds, on average the molecules hit the walls of the container with more force.

5) None of the Above

(1) (Chapter 12)

Ch 12.6 – Kinetic theory of gases

10. What is the average kinetic energy of an N_2 molecule confined in 3.1 L at 1.0 atm and 25°C?


1) $5.71 \times 10^3 \text{ J}$ 2) $9.48 \times 10^3 \text{ J}$ 3) $6.17 \times 10^{-21} \text{ J}$ 4) $5.71 \times 10^{-21} \text{ J}$ 5) $3.21 \times 10^{-21} \text{ J}$

(3) $\overline{K.E.} = \frac{1}{2} m \overline{u^2}$ (per molecule) $\overline{u^2} = \frac{3RT}{M}$

$$\overline{K.E.} = \frac{1}{2} m \frac{3RT}{M} \left(\frac{M N_o^{-1}}{m} \right) = \frac{3}{2} \frac{RT}{N_o} = \frac{3}{2} \frac{(8.314 \text{ J K}^{-1} \text{ mol}^{-1})(25 + 273) \text{ K}}{(6.022 \times 10^{23} \text{ mol}^{-1})} = 6.17 \times 10^{-21} \text{ J}$$

Ch 12.6 – Kinetic theory of gases

Consider the molecular orbital energy diagram shown at right.



11. The energy level denoted “b” refers to:
- 1) a bonding molecular orbital
 - 2) an antibonding molecular orbital
 - 3) a nonbonding molecular orbital
 - 4) an atomic orbital

(2) (OWL question)

Ch 10.3 – basic concepts of molecular orbitals

12. The electrons in the orbital represented by energy level “b”:
- 1) are distributed more toward X
 - 2) are distributed more toward Y
 - 3) are equally distributed between X and Y

(2) (OWL question)

Ch 10, but also 8, 9 – concepts of electronegativities and energy. Covered in class.

13. The molecule XY is the diatomic $(\text{He-H})^+$. What is its bond order?
- 1) 0.0
 - 2) 0.5
 - 3) 1.0
 - 4) 1.5
 - 5) 2.0

(3)

Ch 10, but also 8, 9 – concepts of electronegativities and energy. Covered in class.

14b. What is the energy of ultraviolet light with frequency 1.36×10^{15} Hz?

- 1) 126 kJ mol^{-1}
- 2) 196 kJ mol^{-1}
- 3) 427 kJ mol^{-1}
- 4) 544 kJ mol^{-1}
- 5) 832 kJ mol^{-1}

(4) $E = h\nu = (6.626 \times 10^{-34} \text{ J s})(1.36 \times 10^{15} \text{ Hz}) \left(\frac{\text{s}^{-1}}{\text{Hz}} \right) (6.022 \times 10^{23} \text{ photons mol}^{-1}) = 544000 \text{ J mol}^{-1}$ (OWL question)

Ch 7.2 - light and energy.

15b. Consider two cases for emission from the hydrogen atom:

Case 1:

Electron goes from $n=4$ to $n=3$

Case 2:

Electron goes from $n=6$ to $n=2$

Compare the energies of the photons emitted:

- 1) $E_{\text{case 1}} > E_{\text{case 2}}$ 2) $E_{\text{case 1}} < E_{\text{case 2}}$ 3) $E_{\text{case 1}} = E_{\text{case 2}}$

$$E_n^{H\text{-atom}} = -\frac{R_H hc}{n^2} \quad E_n^{H\text{-atom}} \propto -\frac{1}{n^2}$$

$$\therefore \Delta E = E_f^{H\text{-atom}} - E_i^{H\text{-atom}} \propto -\frac{1}{n_f^2} - \left(-\frac{1}{n_i^2}\right) = \frac{1}{n_i^2} - \left(\frac{1}{n_f^2}\right)$$

$$\Delta E_{\text{case 1}} \propto \frac{1}{3^2} - \left(\frac{1}{4^2}\right) = 20.6 \quad \Delta E_{\text{case 2}} \propto \frac{1}{2^2} - \left(\frac{1}{6^2}\right) = 4.5$$

(1)


Ch 7.3 – hydrogen atom and Rydberg.

16b. Consider the energy vs temperature diagram at right, describing the transitions of water from ice to steam:

The segment labeled (d) is described best with which parameter below:

- 1) $\Delta H^\circ_{\text{fus}}$ 2) $\Delta H^\circ_{\text{vap}}$ 3) C_{ice}
 4) C_{liquid} 5) C_{steam}

(2) boiling



Ch 6.x – phase changes and heat capacities.

17b. The following information is given for bismuth, Bi, at 1atm:

$$\text{boiling pt} = 1627^\circ\text{C} \quad H_{\text{vap}}^{1627^\circ\text{C}, 1\text{atm}} = 172 \text{ kJ mol}^{-1} \quad C_{\text{liquid Bi}} = 0.151 \text{ J g}^{-1} \text{ K}^{-1}$$

$$\text{melting pt} = 271^\circ\text{C} \quad H_{\text{fus}}^{271^\circ\text{C}, 1\text{atm}} = 11.0 \text{ kJ mol}^{-1} \quad C_{\text{solid Bi}} = 0.126 \text{ J g}^{-1} \text{ K}^{-1}$$

At a pressure of 1 atm, what amount of heat is needed to melt a 35.2 g sample of solid bismuth at its normal melting point of 271 °C?

- 1) 4.21 kJ 2) 13.8 kJ 3) 0.561 kJ 4) 9.67 kJ 5) 1.85 kJ

$$(5) \quad q = nH_{\text{fus}}^{271^\circ\text{C}, 1\text{atm}} = \frac{m}{M} H_{\text{fus}}^{271^\circ\text{C}, 1\text{atm}} = \left(35.2 \frac{\text{mol}}{209.0\text{g}} \right) (11.0 \text{ kJ mol}^{-1}) = 1.85 \text{ kJ}$$

Ch 6.x – phase changes and heat capacities.

18b. At a pressure of 1 atm, what amount of heat is needed to take a 35.2 g sample of bismuth from 200°C to 400°C?

- 1) 2.85 kJ 2) 15.4 kJ 3) 32.6 kJ 4) 9.67 kJ 5) 14.3 kJ

(1)

$$\begin{aligned} q &= \frac{m}{M} C_{\text{solid Bi}} + \frac{m}{M} H_{\text{fus}}^{271^\circ\text{C}, 1\text{atm}} + \frac{m}{M} C_{\text{liquid Bi}} \\ &= [0.126 \text{ J g}^{-1} \text{ K}^{-1} (35.2\text{g})(271 - 200)\text{K}] \left(\frac{\text{kJ}}{10^3 \text{ J}} \right) + 1.85 \text{ kJ} + [0.151 \text{ J g}^{-1} \text{ K}^{-1} (35.2\text{g})(400 - 271)\text{K}] \left(\frac{\text{kJ}}{10^3 \text{ J}} \right) \\ &= (0.315 + 1.85 + 0.686) \text{ kJ} = 2.85 \text{ kJ} \end{aligned}$$

Ch 6.x – phase changes and heat capacities.

19b. Which ion has the smallest radius?

- 1) K⁺ 2) Ca²⁺ 3) P³⁻ 4) S²⁻ 5) all the same

(2) – all are isoelectronic. Ca²⁺ has the largest nuclear charge, therefore attracts its electrons the most (OWL 8-12c)

Ch 8.x – ionic radii trends

20b. Consider the following samples:

- a) 0.531 moles of CH₄ in a 6.18 L container at a temperature of 308K
 b) 0.281 moles of CH₄ in a 2.77 L container at a temperature of 388K
 c) 0.569 moles of CH₄ in a 1.42 L container at a temperature of 453K
 d) 0.212 moles of CH₄ in a 5.95 L container at a temperature of 298K

Which has the lowest average molecular speed?

- 1) a 2) b 3) c 4) d 5) all the same

$$(4) \quad \sqrt{u^2} = \sqrt{\frac{3RT}{M}} \quad \text{M all the same; lowest T, lowest rms speed}$$

Ch 12.6 – kinetic theory, rms speed and molar mass.

21b. HNO₃ is (data at the front of the exam provide a clue):

- 1) a strong base 2) a weak acid 3) a weak base
4) a strong acid 5) none of the above

(4) Chapter 5

Ch 5.3 – Acids, but also solubility

22b. Reactions in water that produce gases tend to be:

- 1) favorable 2) ugly 3) unfavorable
4) exothermic 5) endothermic

(1) Chapter 5

Ch 5.5 – Gas forming rxns, but also Ch 6 concepts

23. Which reaction below is a redox reaction?

- 1) NaOH (aq) + HNO₃ (aq) → NaNO₃ (aq) + H₂O (l)
2) Na₂CO₃ (aq) + 2 HClO₄ (aq) → CO₂ (g) + H₂O (l) + 2NaClO₄
3) CdCl₂ (aq) + Na₂S (aq) → CdS (s) + 2 NaCl (aq)
4) Zn(OH)₂ (s) + H₂SO₄ (aq) → ZnSO₄ (aq) + 2 H₂O (l)
5) None of the above

(5) Look at redox changes – there are none. Chapt 5 inspired by book

Ch 5.7 – Redox

24. The net ionic equation for the reaction of zinc sulfate and sodium hydroxide is:

- 1) Zn²⁺ (aq) + 2 OH⁻ (aq) → Zn(OH)₂ (s) + Na₂SO₄ (aq)
2) ZnSO₄ (aq) + 2 NaOH (aq) → Zn(OH)₂ (aq) + Na₂SO₄ (aq)
3) Zn²⁺ (aq) + 2 OH⁻ (aq) → Zn(OH)₂ (s)
4) Zn²⁺ (aq) + 2 OH⁻ (aq) → Zn(OH)₂ (aq)
5) No *net* reaction occurs

(3) hydroxide salts are generally insoluble (OWL 5-2c)

Ch 5.2 – Precipitation rxns

25b. Which element has the lowest ionization energy?

- 1) In 2) Ga 3) Tl 4) B 5) all the same

(3) – IE trends (OWL 8-9b)

Ch 8.x – ionization energy trends


26b. Draw the Lewis structure for CO^{2-} . What is the hybridization on oxygen?

- 1) sp^3d 2) sp^4 3) sp^3 4) sp^2 5) sp

**Ch 10.x and 9 – Lewis structures and hybridization**

27b. Draw the Lewis structure for XeOF_4 (Xe is the central atom). What is the hybridization on Xe?

- 1) sp^3d^2 2) sp^3d^3 3) sp^3d 4) sp^3 5) sp^2

**Ch 10.2 & 9.7 – Hybridization**

28b. The molecule XeOF_4 is:


- 1) polar 2) nonpolar 3) can't tell

**Ch 9.9 – Polarity**

29b. The correct molecular formula for the molecule at right is:

- 1) $\text{C}_2\text{O}_2\text{H}_4$ 2) CO_2H_4 3) C_2OH_4 4) $\text{C}_2\text{O}_2\text{H}_3$

(3)

**Ch 3 – molecular formulas**

30b. A specific isotope of an ion from a given element has 8 protons, 7 neutrons, and 10 electrons. The ion is:

- 1) O^{2-} 2) Ne^{3-} 3) P^{3-} 4) N^{3-} 5) Mn^{3+}

(1) (from an OWL question 3-3c)

Ch 2.3 – atomic composition

31b. What is the formula of the ionic compound formed in the reaction of elemental **Mg** and O_2 ?

- 1) MgO 2) Mg_2O 3) Mg_2O_3 4) Mg_3O_2 5) MgO_2



Ch 3.3 – ionic compounds32b. What is the (mass) percent composition of C in C_4H_8 ?

- 1) 85.6% 2) 14.4% 3) 50.0% 4) 88.3% 5) 11.7%

Mass of C in 1 mol of the compound: $(4 \text{ mol})(12.01 \text{ g mol}^{-1}) = 48.04 \text{ g}$

Mass of 1 mol of the compound:

$$(1 \text{ mol}) \left[4(12.011 \text{ g mol}^{-1}) + 8(1.0079 \text{ g mol}^{-1}) \right] = 56.11 \text{ g}$$

(1) Percent composition: $\frac{48.04 \text{ g C}}{56.11 \text{ g } C_4H_8} 100\% = 85.6\%$ (OWL question)

Ch 3.6 – percent composition33b. What is the wavelength of ultraviolet light with frequency $1.18 \times 10^{15} \text{ Hz}$?

- 1) 209 nm 2) 254 nm 3) 280 nm 4) 190 nm 5) 350 nm

(2) $\lambda = \left(\frac{2.9998 \times 10^8 \text{ m}}{s} \right) \left(\frac{1}{1.18 \times 10^{15} \text{ Hz}} \right) \left(\frac{\text{Hz}}{s^{-1}} \right) \left(\frac{10^9 \text{ nm}}{m} \right) = 254 \text{ nm}$ (OWL question)


Ch 7.1 – wavelength & frequency34b. What is the maximum number of orbitals that can be identified by the set of quantum numbers $n=+5$ $l=+3$?

- 1) 2 2) 3 3) 5 4) 6 5) 7

(5) for $l = 3$, one can have $m_l = -3, -2, -1, 0, +1, +2, +3$ (7 orbitals)

Ch 7.5 – quantum numbers35b. Consider the molecule ClF_4^- How many lone pairs are on the central atom?

- 1) 1 2) 2 3) 3 4) 6 5) 0

**Ch 9.6 – octet rule beyond 2nd row**

36b. Light is given off by a sodium or mercury containing street light when the atoms are excited. The light you see arises for which of the following reasons?


- 1) Electrons are moving from a given energy level to one of lower n
 2) Electrons are moving from a given energy level to one of higher n
 3) Electrons are being removed from the atom, thereby creating a metal cation

(1) (end of chapter question)

Ch 7.3 – atomic energy levels

37b. Consider the molecule ClF_4^- What is the electron pair geometry?

- 1) Trigonal bipyramidal 2) Octahedral 3) linear
4) Trigonal planer 5) Tetrahedral

**Ch 9.7 – electron pair geometry**

38b. Which of the following has the highest affinity for electrons?

- 1) B 2) N 3) As 4) P 5) Ge

(2) (OWL 8-11)

Ch 8.6 – electron affinity

39b. In ionizing elemental lithium to Li^+ , from which orbital is an electron removed?

- 1) 1s 2) 2s 3) 3s 4) 2p 5) 3p

(2) (OWL 8-11)

Ch 8.x – electron configuration and ionization

40b. In the molecule **formaldehyde** CH_2O , what is the approximate HCO bond angle?

- 1) 180° 2) 90° 3) 109° 4) 120° 5) 60°

**Ch 9.7 – molecular geometry**

41. If you completely react 0.678 g of iodine (I_2), what mass of NI_3 can be produced?

- 1) 0.276 g 2) 0.678 g 3) 0.226 g 4) 0.876 g 5) 0.351 g

$$M_{\text{I}_2} = 2(126.9 \text{ g mol}^{-1}) = 253.8 \text{ g mol}^{-1} \quad n_{\text{I}_2} = \frac{0.678 \text{ g}}{253.8 \text{ g mol}^{-1}} = 2.67 \times 10^{-3} \text{ mol}$$


(5)

$$n_{\text{NI}_3} = \frac{1}{3} n_{\text{I}_2} = 8.90 \times 10^{-4} \text{ mol} \quad M_{\text{NI}_3} = 14.01 + 3(126.9 \text{ g mol}^{-1}) = 394.7 \text{ g mol}^{-1}$$

$$m_{\text{NI}_3} = n_{\text{NI}_3} (M_{\text{NI}_3}) = (8.90 \times 10^{-4} \text{ mol})(394.7 \text{ g mol}^{-1}) = 0.351 \text{ g}$$

Ch 3 - Stoichiometry

42. Nitrogen triiodide (NI_3) is unstable, reacting to form N_2 (g) and I_2 (g), and evolving heat.



Spontaneous decomposition of 0.5 g of NI_3 (s) produces what volume of gas at 200°C and 1 atm pressure?

- 1) 28.7 L 2) 0.197 L 3) 0.098 L 4) 14.4 L 5) 0.731 L

$$M = (14.0 + 3(126.9)) \text{ g mol}^{-1} = 394.7 \text{ g mol}^{-1}$$

$$(3) \quad n_{\text{NI}_3} = \frac{0.5 \text{ g}}{394.7 \text{ g mol}^{-1}} = 1.27 \times 10^{-3} \text{ mol}$$

$$n_{\text{gas}} = n_{\text{N}_2} + n_{\text{I}_2} = \frac{1}{2} n_{\text{NI}_3} + \frac{3}{2} n_{\text{NI}_3} = 2 n_{\text{NI}_3} = 2.53 \times 10^{-3} \text{ mol}$$

$$V = \frac{nRT}{P} = \frac{(2.53 \times 10^{-3} \text{ mol})(0.082057 \text{ L atm K}^{-1} \text{ mol}^{-1})(200 + 273) \text{ K}}{1 \text{ atm}} = 0.0983 \text{ L}$$

Ch 3 & 12 – Stoichiometry and gases

43. Using the Table of Bond Dissociation Energies at the front of the exam, predict ΔH° for the spontaneous decomposition of nitrogen triiodide above.

- 1) -256 kJ mol^{-1} 2) -927 kJ mol^{-1} 3) -384 kJ mol^{-1}
 4) -35 kJ mol^{-1} 5) $+927 \text{ kJ mol}^{-1}$


$$(3) \quad \Delta H^\circ = \sum D_{(\text{Bonds broken})} - \sum D_{(\text{Bonds formed})}$$

$$\Delta H^\circ = \{[6(169)] - [945 + 3(151)]\} \text{ kJ} = (1014 - 1398) \text{ kJ} = -384 \text{ kJ}$$

Ch 9.10 – Bond properties

44. What is the molecular geometry of nitrogen triiodide?

- 1) tetrahedral 2) square planar 3) trigonal planar
 4) octahedral 5) trigonal pyramidal

**Ch 9.7 – Molecular Shapes**



45. What is the hybridization on N in nitrogen triiodide?

- 1) sp 2) sp^2 3) sp^3 4) sp^4 5) $sp^3 d$

(3)

Ch 10.2 – Orbital hybridization


46. Which do you expect to have the longest bond length?



5) can't tell

(1)**Ch 8.6 – Atomic properties/trends**

47. In class, we saw the following reaction (unbalanced).



In the correctly balanced reaction, what is the stoichiometry coefficient preceding Br_2 (all coefficients should be integral)?

- 1) 1 2) 2 3) 3 4) 4 5) 6



Ch 4 (but everywhere!) – Balancing reactions

48. In the reaction above of aluminum and bromine, which is the oxidizing agent?

- 1) Al (s) 2) Br₂ (l)



Ch 5.7 - Redox reactions

49. What is the electron pair geometry in AlBr_3 ?

- 1) tetrahedral 2) square planar 3) trigonal planar
4) octahedral 5) trigonal pyramidal

(3)

Ch 9.7 – Molecular Shapes

50. What is the catalog number for this class?

- 1) 123 2) 111 3) 86 4) 345 5) 68.6 g

(2)