
Proliferating Cell Nuclear Antigen


PCNA

- Discovered as an antigen in the autoimmune disease systemic lupus erythematosis
- Co-localized with incorporated bromodeoxyuridine in Sphase
- 1987 Prelich et al. provided proof of PCNA's role in DNA replication
- Subsequent research = processivity factor
 - -PCNA ~10nt incorporation
 - +PCNA ~thousands



Courtesy of John Kuriyan, The Rockefeller University.

Known Binding Partners




www.lbl.gov

DNA * Pol δ and ε Replication Factor C (RFC) FEN1 (flap) endonuclease) ✤ P21 WAF1/CIP1 (CDK) inhibitor) DNA Ligase * XP-G UNG2 MSH2, MLH1 Et cetera


Functions

 Docking partner
DNA polymerase accessory factor
Role in Okazaki Fragment joining
MMR
Chromatin Reorganization
Cell cycle control



PCNA/RFC Interactions

- RFC is the clamp loader
- Head to tail interactions through hydrophobic residues
- Ring must open to accept DNA ATP hydrolysis
- C-side loop (asp41-his44) and c-term tail (lys254-glu256)



PCNA Points of Interests

- Homologous to beta subunit in prokaryotes, similar structure/function, virtually no sequence identity
- Down regulation of PCNA is a marker for some proliferative diseases
- Various malignant tumors are related to expression levels
- Antisense experiments have shown that inhibition can halt certain cancers
- Under proliferative cellular conditions removal of PCNA appears to trigger apoptotic activity
- Many binding partners have a conserved region, -Q--I/L--FF-
- If yours doesn't work right, you are in trouble!

References

- Kontopidis, f. et al., Structural and biochemical studies of human proliferating cell nuclear antigen complexes provide a rationale for cyclin association and inhibitor design. *PNAS*, 102,1871-1876 (2005)
- Prelich, G. et al., The cell-cycle regulated proliferating cell nuclear antigen is required for SV40 DNA replication in vitro. *Nature*, 326, 471-475 (1987)
- Jonsson, Z. O., et al., Regulation of DNA replication and repair proteins through interaction with the front side of proliferating cell nuclear antigen. EMOB J., 17, 2412-2415 (1998)
- Podust, V.N., et al., DNA polymerase delta holoenzyme: action on single-stranded DNA and on double-stranded DNA in the presence of replicative DNA helicases. *Biochemistry*, 34, 5003-5010 (1995)
- Piard, K., et al., Single point mutations located outside the inter-momomer domains abolish trimerization of Schizosaccharomyces pombe PCNA. *Nucleic Acids Research*, 26, 2598-2605 (1998)
- Gary, R. et al., Proliferating Cell Nuclear Antigen Facilitates Excision in Long-patch Base Excision Repair. J. Biological Chemistry, 274, 4354-4363 (1999)
- <www.bioscience.org/1999/v4/d/tsurimot/fulltest.htm>