




Adenosine





















The "Two-Fer" Principle

For hydrogen-bonding in biology, whenever possible, try to get "two for the price of one"

This is more than twice as good

as this



Bringing the two side chains together requires an entropic price

Forming the second H-bond in an already restricted pair does not require paying the entropy price a second time

The "Two-Fer" Principle

For hydrogen-bonding in biology, whenever possible, try to get "two for the price of one"

This is more than twice as good


as this


.o. I. O. I. O.


There is also an electronic argument


Think about alternate resonance forms for the amido group


Resonance Forms






























Good base pairing

Base Pairing


(Donors matched to Acceptors)



Good base pairing Watson-Crick facing

Base Pairing

(Donors matched to Acceptors)




Good base pairing Watson-Crick facing but Anti-Watson-Crick orientation


Good base pairing


Base Pairing

(Donors matched to Acceptors)


Good base pairing WC-Hoogsteen facing








Bad Base Pairing (Donors to Acceptors with terrible angles)







T-Z T


Wild (but good) Base Pairing

Burial of hydrophobic surface drives helix formation (hydrophobic core / stacking interactions)






Burial of hydrophobic surface drives helix formation (hydrophobic core / stacking interactions)




Flat faces are nonpolar


Burial of hydrophobic surface drives helix formation (hydrophobic core / stacking interactions)




Edges are very polar (can H-bond) Flat faces are nonpolar


Other chemical constraints







Furanose Sugar Ring












Furanose Sugar Ring



Furanose Sugar Ring





Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings

TABLE 6-2 A Comparison of the Structural Properties of A, B, and Z DNAs as Derived from Single-Crystal X-Ray Analysis

		indix i ypo	
	Α	В	Z
Overall proportions	Short and broad	Longer and thinner	Elongated and slim
Rise per base pair	2.3 Å	3.32 Å	3.8 Å
Helix-packing diameter	25.5 Å	23.7 Å	18.4 Å
Helix rotation sense	Right-handed	Right-handed	Left-handed
Base pairs per helix repeat	_	_	22
Base pairs per turn of helix	~11	~10	12
Rotation per base pair	33.6°	35.9°	-60° per 2 bp
Pitch per turn of helix	24.6 Å	33.2 Å	45.6 Å
Tilt of base normals to helix axis	+19°	-1.2°	-9°
Base-pair mean propeller twist	+18°	+16°	~0°
Helix axis location	Major groove	Through base pairs	Minor groove
Major-groove proportions	Extremely narrow but	Wide and of intermediate	Flattened out on helix
	very deep	depth	surface
Minor-groove proportions	Very broad but shallow	Narrow and of intermediate	Extremely narrow but
		depth	very deep
Glycosyl-bond conformation	anti	anti	anti at C, syn at G

Source: Adapted from Dickerson R. E. et al. 1982. CSHSQB 47: 14. Copyright @ 1982 Cold Spring Harbor Laboratory Press. Used with permission.

Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings

Nature 430, 377-380 (15 July 2004) | doi: 10.1038/nature02692


Replication by human DNA polymerase-, occurs by

and Aneel K. Aggarwal Deepak T. Nair¹, Robert E. Johnson², Satya Prakash², Louise Prakash² Hoogsteen base-pairing

error-prone) DNA polymerase uses shows that an error correcting (and A crystal structure from from July 2004 recognize the incoming substrate dNTP. Hoogsteen-WC base pairing to

had been suggesting such a pairing. This supports biochemical studies that

This polymerase is good at bypassing lesions in the DNA.



Letters to Nature

Nature 430, 377-380 (15 July 2004) | doi: 10.1038/nature02692

Replication by human DNA polymerase-, occurs by Hoogsteen base-pairing

Deepak T. Nair¹, Robert E. Johnson², Satya Prakash², Louise Prakash² and Aneel K. Aggarwal¹


A crystal structure from from July 2004 shows that an error correcting (and error-prone) DNA polymerase uses Hoogsteen-WC base pairing to recognize the incoming substrate dNTP.

This supports biochemical studies that had been suggesting such a pairing.

This polymerase is good at bypassing lesions in the DNA.

Hoogsteen vindicated


But... perhaps not...





Why is Watson-Crick so good?


Why is Watson-Crick so good?


Why is Watson-Crick so good?




















Why is the major groove so good?

Major Groove


Minor Groove

Why is the major groove so good?

Major Groove

Minor Groove

s me major groove so go





Why is the major groove so good?

Major Groove


Minor Groove


Ribosome






as Benjamin Cummings






Supercoiling




RNA Splicing



Copyright @ 2004 Pearson Education, Inc., publishing as Benjamin Cummings

Translation



Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings